

**ARIZONA GAME AND FISH DEPARTMENT
HERITAGE DATA MANAGEMENT SYSTEM**

Animal Abstract

Element Code: AFCNC05022

Data Sensitivity: No

CLASSIFICATION, NOMENCLATURE, DESCRIPTION, RANGE

NAME: *Poeciliopsis occidentalis sonoriensis*
COMMON NAME: Yaqui topminnow
SYNONYMS: *Girardinus sonoriensis*, *Poeciliopsis sonoriensis*
FAMILY: Poeciliidae

AUTHOR, PLACE OF PUBLICATION: C. Girard, Philadelphia. 1859.

TYPE LOCALITY: Rio San Bernadino, Sonora.

TYPE SPECIMEN: Unknown

TAXONOMIC UNIQUENESS: Two species in the genus in North America. One species occurs in Arizona which has two subspecies, *P.o. occidentalis* and *P.o. sonoriensis*.

DESCRIPTION: "Dorsal profile slightly curved, body somewhat elongated. Caudal fin rounded to almost square. Gonopodium of male elongated, reaching past snout when in copulatory position. Males small, rarely more than 25.0 mm (0.98 in.) standard length; females larger, sometimes 50.0 mm (1.97 in.) or more, usually 30.0 to 45.0 mm (1.18 to 1.77 in.), standard length" (Minckley 1973).

"Body tan to olivaceous, darker above, yellowish to whitened below, but generally lacking dark spots on venter. Scales on dorsum outlined, often diffusely; lateral band developed posterior to pelvic-fin insertions, obsolete anteriorly (or diffuse). Breeding males blackened as in *P.o. occidentalis*, somewhat less intense. Females lacking a darkened peritroct" (Minckley 1973).

AIDS TO IDENTIFICATION: Female topminnow may be distinguished from mosquitofish (*Gambusia affinis*) by lack of dark spots on caudal fin and lack of dark sub-orbital teardrop-shaped mark; origin of dorsal and anal fin vertically inline, perpendicular to horizontal axis of fish, in mosquitofish origin of dorsal fin posterior to origin of anal fin. Male topminnow in breeding condition may or may not become dark black, male mosquitofish never do; male topminnow gonopodium, when extended forward in copulatory position, extends very near to or past snout, male mosquitofish does not. Topminnow have weak, spatulate teeth whereas mosquitofish have strong conically shaped teeth, distinguishable only with a microscope. Female topminnow are generally larger than males.

P.o. sonoriensis can be distinguished from *P.o. occidentalis* by a superior mouth, longer snout, and the lateral bands of the female, which rarely develop before the bases of the pelvic fins (Minckley 1973).

ILLUSTRATIONS: B&W photo (Minckley 1973:202)

TOTAL RANGE: Limited to the Rio Yaqui basin.

RANGE WITHIN ARIZONA: Limited to the portion of the Rio Yaqui basin in the San Bernardino Wildlife Refuge, Arizona.

SPECIES BIOLOGY AND POPULATION TRENDS

BIOLOGY: The topminnow has been found to be able to adjust to wide variations in water quality and appears to be adapted to withstand periodic flash flooding (Galat 1988). Topminnow can withstand water temperatures from near freezing to 32.2-37.7 °C (90-100 F). Long breeding seasons and high fecundity rates have allowed topminnow to rapidly expand their population when habitat is available. Topminnow typically live about one year or a successful breeding season. Actual lifespan is determined by the age at which they reach sexual maturity which may only be a few weeks.

REPRODUCTION: The Yaqui topminnow is a livebearer whose breeding season may last from January through August. Males become dark black and aggressively defend individual territories which females enter when ready to spawn. The male places spermatophores into the female by means of gonopodia. Less dominant males who are unable to defend a territory successfully may attempt inconspicuous breeding by using an extra long gonopodium.

Brood size can vary from 6 to 49 young and gestation lasts 12 to 15 days (Minckley 1973). Females often carry two broods at the same time, each brood at a different stage of development (Minckley 1973). Rinne and Minckley (1991) point out that female has the ability to store sperm, allowing her to fertilize eggs for the duration of her short life from a single copulation.

FOOD HABITS: Omnivorous; detritus, algae, and aquatic invertebrates. Food choice is usually dependent upon a combination of food availability and the topminnow's size.

HABITAT: Lowland and some upland streams of desert and grasslands, and margins of large, lowland rivers. Typical inhabitant of vegetated springs, brooks, and margins and backwaters of larger bodies of water (Minckley 1973). "...topminnows live near the surface in shallow water and are often associated with aquatic vegetation or other cover" (Rinne and Minckley 1991).

ELEVATION: Below 1,500 m (4,920 ft.). According to W. Minckley (AGFD Native Fish Diversity Review 1995), they do not go above 1,300 m (4,265 ft.). Based on records in the Heritage Data Management System (HDMS), elevation ranges from 3,730 - 4,620 ft (1138 - 1409 m) (AGFD, unpublished data accessed 2001).

PLANT COMMUNITY: Cottonwood/willow or burrobrush/weep willow terrestrial riparian communities, in association with aquatic plants such as green algae, *Nasturtium*, *Chara*, and

POPULATION TRENDS: Yaqui topminnow are currently limited to waterways of the San Bernardino National Wildlife Refuge. Loss of habitat and the exotic mosquitofish, *Gambusia affinis*, are thought to be the main threats posed against the subspecies.

SPECIES PROTECTION AND CONSERVATION

ENDANGERED SPECIES ACT STATUS: LE (USDI, FWS 1967)
STATE STATUS: WC (AGFD, WSCA in prep)
[State Endangered AGFD, TNW 1988]
OTHER STATUS: No Forest Service Status (USDA, FS Region 3 1999)
[Forest Service Sensitive, USDA, FS Region 3 1988]
Full Species Listed Threatened (Secretaría de Medio Ambiente 2000)
[Full Species Listed Threatened Secretaría de Desarrollo Social 1994]

MANAGEMENT FACTORS: Water development and diversion, aquifer pumping, reduction in stream flows, drought, introduction of exotic species, and erosion from overgrazing are serious threats to the survival of the Yaqui topminnow. Mosquitofish have caused the elimination of three introduced Yaqui topminnow sites (Bagley et al. 1991). Management needs: protect San Bernardino aquifers, and Leslie Creek and San Bernardino Creek watersheds to ensure adequate perennial flow; identify priority management waters; ameliorate effects of nonnative fishes in management waters; re-establish self-sustaining populations in San Bernardino and Leslie Canyon NWR habitats; stabilize and protect populations in Mexico (AGFD 1996 in prep).

PROTECTIVE MEASURES TAKEN: The Nature Conservancy purchased the San Bernardino Ranch in 1980. The land was then sold to the U.S. Fish and Wildlife Service and the area was established as a national wildlife refuge. In 1984 the Sonoran Topminnow Recovery Plan was drafted.

SUGGESTED PROJECTS: As identified in the Recovery Plan (1983): "1. Maintain, protect, and enhance existing populations and their habitats. 2. Continue surveying waters in the Gila River drainage and the United States portion of the Yaqui River drainage for undiscovered populations of topminnow. 3. Maintain stocks of both Gila and Yaqui topminnow at Dexter National Fish Hatchery. 4. Reintroduce Gila and Yaqui topminnow into suitable sites within the United States portion of their historic ranges. 5. Initiate and support further studies of the Gila and Yaqui topminnow. 6. Enforce all State and Federal laws protecting topminnow populations and their habitat. 7. Develop public support through an information and education program." Work with Mexican authorities to regulate groundwater removal from the San Bernardino valley aquifer.

LAND MANAGEMENT/OWNERSHIP: U.S. Fish and Wildlife Service - San Bernardino National Wildlife Refuge; Johnson Historical Museum; Private.

SOURCES OF FURTHER INFORMATION

LITERATURE CITATIONS:

- Arizona Game and Fish Department. 1988. Threatened Native Wildlife in Arizona. p.31.
- Arizona Game and Fish Department. In prep. Wildlife of special concern in Arizona. Arizona Game and Fish Department Publication. Phoenix, Arizona. 32 pp.
- Arizona Game and Fish Department Native Fish Diversity Review. 1995. Tempe, Arizona.
- Bagley, B.E. et. al. 1991. Status of the Sonoran Topminnow (*Poeciliopsis occidentalis*) and Desert Pupfish (*Cyprinodon macularius*) in Arizona. Special Report on Project E5-2, Job 9 Title of the Endangered Species Act. Arizona Game and Fish Department, Phoenix, Arizona. pp. 64.
- Galat, D.L. and B. Robertson. 1988. Interactions between *Poeciliopsis occidentalis sonoriensis* and *Gambusia affinis* (Pisces:Poeciliidae) in the Rio Yaqui drainage, Arizona. pp. 66.
- Hedrick, P.W. 2000. Final Report on MHC Variation in Topminnows and resistance to Parasites. Final Report, AGFD Heritage Grant # I97011.
- Hedrick, P.W. 2000. Microsatellite and MHC variation to distinguish natural lineages of the Sonoran Topminnow. Final Report, AGFD Heritage Grant # I98002.
- Hendrickson, D.A., W.L. Minckley, R.R. Miller, D.J. Siebert, and P.H. Minckley. 1980. Fishes of the Rio Yaqui basin, Mexico and United States. Journal of the Arizona-Nevada Academy of Science, 15(3):80.
- Minckley, W.L. 1973. Fishes of Arizona. Arizona Game and Fish Department, Phoenix. pp. 198, 202-203.
- Page, L.M. and B.M. Burr. 1991. A field guide to freshwater fishes: North America, north of Mexico. Houghton Mifflin Co., Boston. pp. 239-240.
- Rinne, J.N. and W.L. Minckley. 1991. Native fishes of arid lands: a dwindling resource of the desert southwest. U.S. Department of Agriculture, Forest Service, General Technical Report RM-206. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado. pp. 26-27.
- Secretaría de Desarrollo Social. 1994. Diario Oficial de la Federacion. p. 52.
- Secretaría de Medio Ambiente. 2000. Diario Oficial de la Federacion, PROY-NOM-059-ECOL-2000. p. 46.
- USDA, Forest Service Region 3. 1988. Regional Forester's Sensitive Species List.
- USDA, Forest Service Region 3. 1999. Regional Forester's Sensitive Species List.
- USDI, Fish and Wildlife Service. 1967. Endangered and Threatened Wildlife and Plants; Review of Vertebrate Wildlife; Notice of Review. Federal Register 32:4001.
- USDI, Fish and Wildlife Service. 1983. Gila and Yaqui Topminnow Recovery Plan. U.S. Fish and Wildlife Service, Albuquerque, New Mexico. pp. 56.

MAJOR KNOWLEDGEABLE INDIVIDUALS:

- Jeff Simms - Bureau of Land Management, Safford District, Safford, Arizona.
- Jerry Stefferud - USDA, Forest Service, Tonto National Forest, Phoenix.

AGFD Animal Abstract

-5-

Poeciliopsis occidentalis sonoriensis

Sally Stefferud - USDI, Fish and Wildlife Service, Ecological Services, Phoenix.

David A. Weedman - Arizona Game and Fish Department, Nongame Branch, Phoenix.

ADDITIONAL INFORMATION:

Revised: 1994-09-15 (SBZ)
1994-09-23 (MHH)
2001-10-18 (SMS)

To the user of this abstract: you may use this entire abstract or any part of it. We do request, however, that if you make use of this abstract in plans, reports, publications, etc. that you credit the Arizona Game and Fish Department. Please use the following citation:

Arizona Game and Fish Department. 2001. *Poeciliopsis occidentalis sonoriensis*.
Unpublished abstract compiled and edited by the Heritage Data Management System,
Arizona Game and Fish Department, Phoenix, AZ. 4 pp.